

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) Publication number:

**0 485 079 A1**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: **91309508.9**

(51) Int. Cl.<sup>5</sup>: **A01N 47/44, //(A01N47/44,  
25:02,25:22)**

(22) Date of filing: **16.10.91**

(30) Priority: **06.11.90 GB 9024133**

(43) Date of publication of application:  
**13.05.92 Bulletin 92/20**

(94) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IT LI LU NL SE**

(71) Applicant: **IMPERIAL CHEMICAL INDUSTRIES  
PLC**  
**Imperial Chemical House, Millbank**  
**London SW1P 3JF(GB)**

(72) Inventor: **Henry, William John**  
**34 Tarmangie Drive**  
**Dollar, Clackmannanshire FK14 7BP(GB)**

(74) Representative: **Pugsley, Roger Graham et al**  
**IMPERIAL CHEMICAL INDUSTRIES PLC Legal**  
**Department: Patents PO Box 6 Bessemer**  
**Road**  
**Welwyn Garden City Herts, AL7 1HD(GB)**

(54) **Aqueous composition.**

(57) A aqueous medium containing a biguanide, particularly a biguanide oligomer, and having a pH of less than 5. The biguanide oligomer may be a poly(hexamethylene biguanide) having up to 15 repeat units. The medium has improved storage stability compared to commercially available systems and is at least as effective as a biocide.

EP 0 485 079 A1

The present invention relates to aqueous compositions and particularly to aqueous compositions containing a biguanide, especially a biguanide oligomer.

Biguanides, particularly biguanide oligomers such as hexamethylene biguanide oligomers having up to 15 repeat units, are commercially available materials which can be used as industrial biocides or disinfectants. These materials are typically available in an aqueous medium at a pH of 5 or greater, typically about pH 5.5, and at a concentration of the active ingredient of about 20% by weight relative to the weight of the total composition. It has been found that after a period of time, haze formation can occur and this is generally undesirable and may be unacceptable in some applications.

Hence, it is desirable to provide an aqueous composition containing a biguanide, particularly a biguanide oligomer, which has an improved resistance to the formation of haze.

According to the present invention there is provided an aqueous composition which contains a biguanide and which has a pH of less than 5.


We have found that the pH of aqueous compositions containing biguanides, particularly such compositions containing biguanide oligomers, is dependent on the technique used to measure the pH. Hitherto we have measured, and reported, the pH of aqueous compositions containing biguanides by the use of a pH meter using a glass electrode. However, we have found that the pH so determined can show an upward drift with time and, furthermore, does not correspond to the pH of the same composition as determined using an indicator such as universal indicator. The reason for this difference in the measured pH is believed to be that the aqueous medium is not a true solution but rather one in which the biguanide is present as a second, finely dispersed, phase in the aqueous medium. It is believed that the heterogeneity of such a two phase system is the cause of misleading results obtained when using a glass electrode. Accordingly, unless indicated to the contrary, the term "pH" as used hereafter is the pH obtained using an indicator appropriate for the pH being determined.

The pH of the aqueous composition may be less than 1, for example 0.1 but is typically not less than 2, and generally is at least 3 and especially is at least 4 and less than 5. A pH in the range at least 4 and less than 5 is conveniently measured using bromo-cresol green as the indicator but it will be appreciated that any indicator may be used provided it is appropriate for measurement in the pH range being determined.

A pH of less than 5, as measured using an appropriate indicator, typically has a numerical value which is about one more than the pH of the same composition as measured using a glass electrode. The aqueous composition of the present invention typically has a pH of not more than 4.8 and especially not more than 4.5. We have found that a pH in the range 4 to 4.5 corresponds to a pH of about 3 to 3.5 when determined using a glass electrode.


We have found that the aqueous compositions of the present invention can be stored for a longer period of time than commercially available products with little haze formation. Furthermore, the aqueous compositions when subjected to a microbiological evaluation of activity show essentially the same activity as the commercially available products and in some tests have shown a somewhat greater activity.

The biguanide which is present in the aqueous composition of the present invention contains at least one biguanide unit of the formula (I):-




Typically the biguanide contains at least two units of the formula (I) which are linked by a bridging group which contains at least one methylene group. The bridging group may include a polymethylene chain which may optionally be interrupted by hetero atoms such as oxygen, sulphur or nitrogen. The bridging group may include one or more cyclic nuclei which may be saturated or unsaturated. It is generally preferred that the bridging group is such that there are at least three, and especially at least four, carbon atoms directly interposed between two adjacent units of the formula (I). In general it is preferred that there are not more than ten carbon atoms, especially not more than eight carbon atoms, interposed between two adjacent units of the formula (I).

The biguanide units may be terminated by any suitable group which may be a hydrocarbyl or substituted hydrocarbyl group or which may be an amine group or an amine hydrochloride group or by a group




If the terminating group is a hydrocarbonyl group this may be an alkyl, cycloalkyl or aryl group or may be a combination thereof as in an aralkyl group. If the terminating group is a substituted hydrocarbonyl group, the substituent can be any substituent which does not have an undesirable adverse effect on the microbial activity of the biguanide compound and typically is a hydrocarbonoxy group, a hydrocarboncarbonyl (that is an acyl) group, an ester (that is an acyloxy) group, a halogen atom or a nitrile group and there may be more than one substituent, for example more than one halogen atom.

A suitable biguanide is a material which contains two units of the formula (I) and in which the units are linked by a polymethylene group, particularly a hexamethylene group. The terminating groups may be 4-chlorophenyl groups, for example as in the compound of formula (II):-




The compound of formula (II) is available as a chlorhexidine salt.

The biguanide may alternatively be a polymeric biguanide, for example a linear polymeric biguanide which has a recurring polymer unit represented by the formula



wherein X and Y may be the same or different and represent bridging groups in which together the total number of carbon atoms directly interposed between the pairs of nitrogen atoms linked by X and Y is at least 9 and not more than 17.


The bridging groups X and Y may consist of polymethylene chains, optionally interrupted by hetero atoms, for example, oxygen, sulphur or nitrogen. X and Y may also incorporate cyclic nuclei which may be saturated or unsaturated, in which case the number of carbon atoms directly interposed between the pairs of nitrogen atoms linked by X and Y is taken as including that segment of the cyclic group, or groups, which is the shortest. Thus, the number of carbon atoms directly interposed between the nitrogen atoms in the group




is 4 and not 8.

The preferred polymeric biguanide for use in the present invention is poly(hexamethylene biguanide), in which X and Y both represent the  $-(CH_2)_6-$  group.

Polymeric biguanides may be prepared by the reaction of a bisdicyanidiamide having the formula




with a diamine  $H_2N-Y-NH_2$ , wherein X and Y have the meanings defined previously herein; or by reaction between a diamine salt of dicyanimide having the formula




with a diamine  $H_2N-Y-NH_2$  wherein X and Y have the meanings defined previously herein. These methods of preparation are described in UK Patent Specifications Nos. 702,268 and 1152243 respectively, and any of the polymeric biguanides described therein may be used as the biguanide component of the aqueous composition of the present invention.

The biguanide polymer chains are terminated either by an amino hydrochloride group or by an




and the terminating groups may be the same or different on each polymer chain.

Typically, the polymeric biguanides are obtained as mixtures of polymers in which the polymer chains are of different lengths, the number of individual biguanide units, i.e.




and



together being from 3 to about 80.

In the case of the preferred poly(hexamethylene biguanide) having the formula (III)



the value of  $n$  is in the range from 4 to 15, the average molecular weight of the polymer mixture being from about 1100 to about 3300.

The biguanides are used as salts with suitable inorganic or organic acids, for example as the hydrochloride salts or the acetate or gluconate.

The preferred aqueous compositions in accordance with the present invention contain a poly-(hexamethylene biguanide) of formula III as the hydrochloride salt and have a pH of at least 4 up to 4.5.

The biguanide may be used to provide microbiological protection in an amount of less than 1% w/w, for example in amounts of less than 1000 ppm, especially less than 250 ppm by weight. However, the aqueous compositions of the present invention are typically used as concentrated solutions which can be diluted to a desired level in use. Thus, the biguanide is typically present in the aqueous composition in an amount of at least 1% w/w. In general, the concentration of the biguanide will be at least 5% w/w. The biguanide may be present at concentrations of 25% w/w or even higher but at such concentrations the problem of haze formation is increased. The biguanide is conveniently used at a concentration of about 20% w/w.

We have found the aqueous compositions of the present invention generally show little, if any, haze on being stored for a period of up to two months at ambient temperature in closed drums whereas a similar aqueous composition having a pH in the range 6.5 to 7.0 (about 5.5 using a glass electrode) shows considerable turbidity after being stored for this period of time.

The pH of the aqueous composition may be adjusted to the desired value using a suitable acid or base, typically by the addition of an acid such as hydrochloric acid.

Biguanides have anti-microbial properties and the aqueous compositions of the present invention may be used in any of the applications for which commercially available biguanide compositions have been used or recommended, for example as preservatives for personal-care products, for the protection of hides and skins, and in disinfectants for use in brewing, food processing and the like.

Thus, as a further aspect of the present invention there is provided a method for inhibiting the growth of micro-organisms on, or in, a medium which comprises treating the medium with an aqueous composition containing a biguanide and as hereinbefore defined. The aqueous composition may contain only the biguanide as the anti-microbial agent.

The aqueous composition can be used in conditions in which micro-organisms grow and cause problems. Systems in which micro-organisms cause problems include liquid, particularly aqueous, systems such as cooling water liquors, paper mill liquors, metal working fluids, geological drilling lubricants, polymer emulsions and surface coating compositions such as paints, varnishes and lacquers and also solid materials such as wood and leather. The aqueous composition of the present invention can be included in such materials to provide an anti-microbial effect. The amount of the aqueous composition is typically sufficient to provide a biguanide concentration in the range from 0.0001 up to 10%, preferably 0.0002 up to 5% and especially 0.0002 to 0.1% by weight of biguanide relative to the weight of the system to which it is added. In many cases, microbial inhibition may be obtained with between 0.0005% and 0.05% by weight of the biguanide.

The biguanides which are present in the aqueous composition of the present invention may be the only antimicrobial compounds or may be used together with further compounds having antimicrobial characteristics. The aqueous composition may contain more than one biguanide compound of general formula I. Alternatively, an aqueous composition of a biguanide compound of pH less than 5 in accordance with the present invention may be used together with one or more known antimicrobial compounds. The use of a mixture of anti-microbial compounds can provide a composition having a broader anti-microbial spectrum and hence one which is more generally effective than the components thereof. The known antimicrobial may be one possessing anti-bacterial, anti-fungal, anti-algal or other antimicrobial characteristic. The mixture of the biguanide with other antimicrobial compounds typically contains from 1 to 99% by weight, and particularly from 40 to 60% by weight, relative to the weight of total antimicrobially active compounds, of the mixture of a biguanide compound.

As examples of known antimicrobial compounds which may be used, together with a biguanide compound there may be mentioned quaternary ammonium compounds such as diethyldodecylbenzyl ammonium chloride; dimethyloctadecyl(dimethylbenzyl)ammonium chloride; dimethyldidecylammonium chloride; dimethyldidodecylammonium chloride; trimethyltetradecylammonium chloride; benzyltrimethyl-  
 5 (C<sub>12</sub>-C<sub>18</sub> alkyl)ammonium chloride; dichlorobenzyltrimethyldodecylammonium chloride; hexadecylpyridinium chloride; hexadecylpyridinium bromide; hexadecyltrimethylammonium bromide; dodecylpyridinium chloride; dodecylpyridinium bisulphate; benzyl-dodecyl-bis(beta-hydroxyethyl)ammonium chloride; dodecylbenzyl-trimethylammonium chloride; benzyltrimethyl(C<sub>12</sub>-C<sub>18</sub> alkyl) ammonium chloride; dodecyltrimethylethyl am-  
 10 monium ethylsulphate; dodecyltrimethyl-(1-naphthylmethyl)ammonium chloride; hexadecyltrimethylbenzyl ammonium chloride; dodecyltrimethylbenzyl ammonium chloride and 1-(3-chloroallyl)-3,5,7-triaza-1-azonia-  
 adamantane chloride; urea derivatives such as 1,3-bis(hydroxymethyl)-5,5-dimethylhydantoin; bis-  
 (hydroxymethyl)urea; tetrakis(hydroxymethyl)acetylene diurea; 1-(hydroxymethyl)-5,5-dimethylhydantoin  
 and imidazolidinyl urea; amino compounds such as 1,3-bis(2-ethyl-hexyl)-5-methyl-5-am-  
 15 inohexahydropyrimidine; hexamethylene tetra amine; 1,3-bis(4-aminophenoxy)propane; and 2-[(  
 hydroxymethyl)- amino]ethanol; imidazole derivatives such as 1-[2-(2,4-dichloro- phenyl)-2-(2-propenyloxy)-  
 ethyl]-1H-imidazole; 2-(methoxycarbonyl- amino)-benzimidazole; nitrile compounds such as 2-bromo-2-  
 bromomethylglutaronitrile, 2-chloro-2-chloromethylglutaronitrile, 2,4,5,6-tetra- chloroisophthalodinitrile;  
 thiocyanate derivatives such as methylene bis thiocyanate; tin compounds or complexes such as tributyltin-  
 20 oxide, chloride, naphthoate, benzoate or 2-hydroxybenzoate; isothiazolin-3-ones such as 4,5-trimethylene-4-  
 isothiazolin-3-one, 2-methyl-4,5-trimethylene-4-isothiazolin-3-one, 2-methylisothiazolin-3-one, 5-chloro-2-  
 methyl-isothiazolin-3-one, benzisothiazolin-3-one and 2-methylbenzisothiazolin-3-one; thiazole derivatives  
 such as 2-(thiocyanomethylthio)-benzthiazole; and mercaptobenzthiazole; nitro compounds such as tris-  
 (hydroxymethyl)nitromethane; 5-bromo-5-nitro-1,3-dioxane and 2-bromo-2-nitropropane-1,3-diol; iodine com-  
 25 pounds such as iodo propynyl butyl carbamate and tri-iodo allyl alcohol; aldehydes and derivatives such as  
 glutaraldehyde (pentanedial), p-chlorophenyl-3-iodopropargyl formaldehyde, and glyoxal; amides such as  
 chloracetamide; N,N-bis(hydroxymethyl)chloracetamide; N-hydroxymethyl-chloracetamide and dithio-2,2-bis-  
 (benzomethyl amide); thiones such as 3,5-dimethyltetrahydro-1,3,5-2H-thiadiazine-2-thione; triazine deriva-  
 tives such as hexahydrotriazine and 1,3,5-tri-(hydroxyethyl)-1,3,5-hexahydrotriazine; oxazolidine and deriva-  
 tives thereof such as bis-oxazolidine; furan and derivatives thereof such as 2,5-dihydro-2,5-dialkoxy-2,5-  
 30 dialkylfuran; carboxylic acids and the salts and esters thereof such as sorbic acid and the salts thereof and  
 4-hydroxybenzoic acid and the salts and esters thereof; phenol and derivatives thereof such as 5-chloro-2-  
 (2,4-dichloro- phenoxy)phenol; thio-bis(4-chlorophenol) and 2-phenylphenol; sulphone derivatives such as  
 diiodomethyl-paratolyl sulphone, 2,3,5,6-tetrachloro-4-(methylsulphonyl) pyridine and hexachlorodimethyl  
 sulphone.

35 Further aspects of the present invention are described in the following illustrative examples.

### Example 1

A commercially available aqueous composition having a pH of 5.3 as measured by a pH meter with a  
 40 glass electrode, and containing 20% w/w of poly(hexamethylene biguanide) having an average number of  
 repeating units in the range 4-7 was found to have a pH in the range 6.5 to 7.0 as determined by Lyphan  
 pH indicator paper (universal indicator paper).

To a sample of the above aqueous composition was added sufficient concentrated (37%) aqueous  
 hydrochloric acid to give a pH of 3.5 as measured by a pH meter with a glass electrode. The pH of this  
 45 acidified material was also determined using bromocresol green indicator and was found to be 4.6 by this  
 method. The pH was determined by the addition of 10 drops of 0.04% bromocresol green indicator solution  
 to 50cm<sup>3</sup> of the acidified material in standard 50cm<sup>3</sup> Nessler tubes which were then introduced into a  
 Nessleriser comparator disc apparatus (BDH Lovibond AF 300 Mark 3).

Samples of the commercially available aqueous composition and of the acidified material were stored in  
 50 25 kg drums for nine months under ambient conditions (temperature 15-25°C). After storage, the samples  
 were visually inspected for turbidity. The commercially available aqueous composition showed slight  
 turbidity whilst the acidified material was assessed as "Perfect Clarity".

Six further pairs of samples were prepared essentially as described and were stored for two months. In  
 all cases, the acidified material was assessed as "Perfect Clarity" whilst five samples of the commercially  
 55 available aqueous composition were assessed as "Turbid" the remaining sample being intermediate  
 between "Slight Turbidity" and "Turbid".

### Examples 2 and 3

The commercially available aqueous composition used in Example 1 was acidified to give samples of pH 4.6 and pH 4.4 (determined using bromocresol green as described). Samples of all three compositions were stored for one month, conditions of storage being as described in Example 1. By visual inspection after one month's storage, the commercially available aqueous composition showed heavy turbidity whilst both acidified materials show perfect clarity.

Three further sets of samples were prepared essentially as described and stored for one month. In all cases, the commercially available aqueous composition was assessed as showing heavy turbidity. One sample of pH 4.6 was assessed as being turbid but all other acidified materials were assessed as showing perfect clarity.

### Example 4

100 cm<sup>3</sup> samples of an aqueous solution containing a range of concentrations of poly(hexamethylene biguanide) were prepared from either the commercially available aqueous composition as used in Example 1 or from an acidified material of pH 4.6 (using bromocresol green indicator). To each sample was added a bacterial inoculum (*Escherichia coli*) to give a bacterial concentration in the liquid medium of about  $1 \times 10^8$  cells/cm<sup>3</sup>. The samples were placed in 250 cm<sup>3</sup> conical flasks, and contained poly(hexamethylene biguanide) at concentrations of 200, 100, 50, 25, 12.5 or 6.25 ppm.

The solutions containing poly(hexamethylene biguanide), together with a solution containing no additive, were incubated at ambient temperature (15°-25°C). After incubation periods of 10 minutes and one hour, the surviving bacteria were determined by the decimal dilution method using nutrient agar.

The results obtained are set out in the following Table.

TABLE

Sample (a)	PHMB Conc. (ppm) (b)	Survivors (cells/cm <sup>3</sup> )	
		10 min	one hour
1	200	<10	<10
1	100	<10	<10
1	50	<10	<10
1	25	<10	<10
1	12.5	$1.0 \times 10^3$	<10
1	6.25	$1.2 \times 10^4$	<10
A	200	<10	<10
A	100	$5 \times 10^1$	<10
A	50	$1.2 \times 10^2$	<10
A	25	$8.6 \times 10^3$	<10
A	12.5	$1.1 \times 10^4$	<10
A	6.25	$3.5 \times 10^5$	<10
-	NIL	$3.0 \times 10^8$	$4.5 \times 10^8$

Notes to Table

(a) 1 is a acidified poly(hexamethylene biguanide)aqueous solution of pH 4.6

A is a commercially available poly(hexamethylene biguanide)aqueous solution of pH 6.5-7.0.

(b) The concentration is that of the active ingredient, PHMB is poly(hexamethylene biguanide).

Examples 5 to 7

To different samples of the commercially available aqueous composition described in Example 1 were added sufficient quantities of concentrated (37%) aqueous hydrochloric acid to give materials having a pH of 2.8, 1.8 and 1.0 as determined by bromocresol green, malachite green and malachite green respectively.

50 cm<sup>3</sup> samples of each of these acidified materials were placed in jacketed vessels and the contents of each vessel were stirred. The contents of the vessels were subjected to a "freeze-thaw" test by cooling to -8°C and heating to 15°C using ethylene glycol as the heat transfer liquid. Heating and cooling were




carried out at a rate of 0.6°C per minute. The temperature of -8°C was maintained for one minute and the temperature of 15°C was maintained for 15 minutes. The "freeze-thaw" cycle was effected a total of 15 times.

At the end of 15 cycles, the samples were removed and examined. All the samples were assessed as "Perfect Clarity".


# Claims

1. An aqueous composition which contains a biguanide and has a pH of at least 0.1 and less than 5.
2. An aqueous composition as claimed in claim 1 which has a pH of at least 4 and less than 5.
3. An aqueous composition as claimed in either claim 1 or claim 2 wherein the biguanide contains at least two units of the formula I



which are linked by a bridging group which contains at least one methylene group.

4. An aqueous composition as claimed in claim 3 wherein the bridging group is a hexamethylene group.
5. An aqueous composition as claimed in claim 4 in which the biguanide is poly(hexamethylene biguanide) having the formula (III)



wherein the value of n is from 4 to 15.

6. An aqueous composition as claimed in claim 5 wherein the poly(hexamethylene biguanide) is in the form of its hydrochloride salt.
7. An aqueous composition as claimed in any one of claims 1 to 6 which contains up to 25% w/w of the biguanide.
8. A method for inhibiting the growth of micro-organisms on, or in, a medium which comprises treating the medium with an aqueous composition as claimed in any one of claims 1 to 7.
9. A method as claimed in claim 8 wherein the aqueous composition is added in an amount to provide a biguanide concentration in the range from 0.0001 up to 10% by weight of biguanide relative to the medium.
10. A method as claimed in either claim 8 or claim 9 wherein the biguanide is used together with further compounds having antimicrobial characteristics.



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number

EP 91 30 9508

### DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	GB-A-2 187 097 (UNILEVER PLC) * page 1, line 15 - page 1, line 57; claims * ---	1-4, 7, 8	A01N47/44 /(A01N47/44, 25:02, 25:22)
X	EP-A-0 110 568 (THE PROCTER & GAMBLE COMPANY) * page 2, line 13 - page 2, line 39 * * page 3, line 25 - page 5, line 2; claims; examples * ---	1-4, 7, 8	
X	EP-A-0 379 256 (COLGATE-PALMOLIVE COMPANY)  * page 3, line 45 - page 3, line 55 * * page 5, line 51 - page 6, line 6; claims; examples * ---	1, 3, 4, 7-10	
X	DE-A-3 723 976 (CIBA-GEIGY AG)  * column 3, line 61 - column 3, line 65 * * column 5, line 27 - column 5, line 28; claims * -----	1, 3, 4, 7-10	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			A01N
Place of search THE HAGUE		Date of completion of the search 10 FEBRUARY 1992	Examiner DONOVAN T. M.
<b>CATEGORY OF CITED DOCUMENTS</b>			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document			